

available at <http://www.tzuchimedjnl.com/>



Tzu Chi Medical Journal

Original Article

Magnesium Sulfate Inhibits Activator Protein-1 Upregulation in Endotoxin-activated Murine Macrophages

Cay-Huyen Chen^{1,2}, Pei-Shan Tsai³, Chun-Jen Huang^{1,2*}

¹Department of Anesthesiology, Buddhist Tzu Chi General Hospital, Taipei Branch, Taipei, Taiwan

²School of Medicine, Tzu Chi University, Hualien, Taiwan

³College of Nursing, Taipei Medical University, Taipei, Taiwan

Article info

Article history:

Received: August 12, 2010

Revised: September 23, 2010

Accepted: October 13, 2010

Keywords:

AP-1

LPS

L-type calcium channels

MgSO₄

Abstract

Objectives: The expression of inflammatory molecules is regulated by the transcription factor activator protein-1 (AP-1), a heterodimeric protein that consists of proteins from various families, including c-Jun and c-Fos. We sought to elucidate whether MgSO₄ regulates the activation of AP-1 in endotoxin-activated RAW264.7 cells, a murine macrophage-like cell line. The possible roles of the L-type calcium channels in this process were also elucidated.

Materials and Methods: RAW264.7 cells were treated with phosphate buffered saline, MgSO₄, lipopolysaccharide (LPS), LPS plus MgSO₄ (20 mM), or LPS plus MgSO₄ plus the L-type calcium channel activator BAY-K8644 (1 μM). After harvesting, expression of AP-1 was evaluated.

Results: LPS induced AP-1 activation based on the fact that the nuclear protein concentrations of AP-1 components, including c-Jun and c-Fos, as well as the AP-1 DNA-binding activity, were significantly increased in LPS-treated RAW264.7 cells. MgSO₄, in contrast, significantly inhibited LPS-induced AP-1 activation in activated RAW264.7 cells. Moreover, the effect of MgSO₄ on AP-1 was reversed by BAY-K8644.

Conclusion: MgSO₄ inhibited AP-1 activation in LPS-treated macrophages and the mechanism may involve the L-type calcium channels. (*Tzu Chi Med J* 2010;22(4):177–183)

*Corresponding author. Department of Anesthesiology, Buddhist Tzu Chi General Hospital, Taipei Branch, 289, Jianguo Road, Sindian, Taipei, Taiwan.
E-mail address: huangcj1112@gmail.com

1. Introduction

Upregulation of inflammatory molecules is essential in mediating the development of systemic inflammatory responses during sepsis (1). Therapies aimed at decreasing inflammatory molecule expression have been shown to be effective in reducing the pathological sequelae of sepsis (2). Transcription factors, including nuclear factor-κB (NF-κB) and activator protein-1 (AP-1),

are essential for the production of endotoxin-induced inflammatory molecules (3,4). AP-1 is a heterodimeric protein that consists of proteins from various families, including c-Jun, c-Fos and others (5).

Magnesium sulfate (MgSO₄) is an anticonvulsant that is used for the treatment of severe preeclampsia and eclampsia (6). MgSO₄ also reduces intraoperative anesthetic requirements and is effective in controlling neuropathic pain (7,8). Moreover, MgSO₄ has

been shown to possess certain anti-inflammation effects. Our previous data (9), in agreement with other investigations (10), confirmed that MgSO_4 significantly suppressed the inflammatory responses in cells treated with endotoxin. MgSO_4 has also been reported to suppress NF- κ B activation (9,10).

However, the effects of MgSO_4 on regulating the activation of AP-1 remain unstudied. To elucidate further, we thus conducted this cellular study based on the hypothesis that MgSO_4 is able to inhibit the activation of AP-1 in endotoxin-activated murine macrophages. In addition, it is known that magnesium is a potent L-type calcium channel inhibitor (11). Our previous results have demonstrated the involvement of L-type calcium channels in mediating the anti-inflammation effects of MgSO_4 (9). Therefore, in this study, we investigated the possible role of the L-type calcium channels with respect to MgSO_4 and the activation of AP-1.

2. Materials and methods

2.1. Cell culture and cell activation protocols

RAW264.7 cells, an immortalized murine macrophage-like cell line, is able to readily express inflammatory molecules upon stimulation by endotoxin (12). Based on this fact, we employed RAW264.7 cells in this investigation. RAW264.7 cells were grown in Dulbecco's modified Eagle's medium (Life Technologies, Grand Island, NY, USA) supplemented with 10% fetal bovine serum and 1% penicillin/streptomycin (Life Technologies) and incubated in a humidified chamber at 37°C using a mixture of 95% air and 5% CO_2 . Confluent RAW264.7 cells were then activated with lipopolysaccharide (LPS, 100 ng/mL, *Escherichia coli* serotype 0127:B8 endotoxin; Sigma-Aldrich, St. Louis, MO, USA) to induce the expression of the various molecules to be investigated, as previously reported (13).

2.2. Experimental protocols

Confluent RAW264.7 cells were randomly allocated into five groups. (1) The PBS group: RAW264.7 cells were treated with 1× phosphate buffered saline (PBS, Life Technologies). (2) The LPS group: RAW264.7 cells were treated with LPS (100 ng/mL; Sigma-Aldrich). (3) The M group: RAW264.7 cells were treated with MgSO_4 (20 mM; Sigma-Aldrich). (4) The LPS+M group: RAW264.7 cells were treated with LPS plus MgSO_4 (20 mM). (5) The LPS+M+K group: RAW264.7 cells were treated with LPS plus MgSO_4 (20 mM) plus a potent L-type calcium channel activator BAY-K8644 (1 μ M; Sigma-Aldrich). The doses of MgSO_4 and BAY-K8644 were determined based on our previous data, which

indicated that MgSO_4 at 20 mM was able to attenuate endotoxin-induced upregulation of inflammatory cytokines in RAW264.7 cells and that BAY-K8644 at 1 μ M was able to reverse the abovementioned effects of MgSO_4 (9). MgSO_4 was administered immediately after LPS. BAY-K8644 was administered 5 minutes prior to LPS and was followed by MgSO_4 . Each group contained 24 culture dishes ($n=24$ in each group). For the analysis of AP-1, six culture dishes from each group were harvested after reacting with LPS for 0, 1, 3, and 5 hours. Similar durations were used for the groups without LPS.

2.3. Nuclear extract preparation and immunoblotting assay

To facilitate assaying the activation of AP-1, including c-Jun and c-Fos, nuclear extracts were prepared as previously reported (13). In brief, cells were washed, scraped, and centrifuged at 1500g for 5 minutes. The cell pellet was resuspended in 5 mL of cell lysis buffer (10 mM HEPES (pH 7.9), 1.5 mM MgCl_2 , 10 mM KCl, and 0.6% NP-40; all chemicals were purchased from Sigma-Aldrich) and centrifuged again at 1500g for 5 minutes. Cells were resuspended again in cell lysis buffer and allowed to swell on ice for 10 minutes followed by homogenization. The homogenate was centrifuged at 3300g for 15 minutes at 4°C. After discarding the supernatant, the pellet was resuspended and homogenized in three volumes of nuclear extraction buffer (20 mM HEPES (pH 7.9), 1.5 mM MgCl_2 , 420 mM NaCl, 0.2 mM EDTA, and 25% glycerol; Sigma-Aldrich). After stirring on ice for 30 minutes and centrifuging at 89,000g for 30 minutes, the supernatant from the nuclear suspension was collected and concentrated in a Microcon 10 concentrator (Millipore Corp., Burlington, MA, USA) by centrifugation at 14,000g for 3 hours at 4°C. The protein concentration of each sample was measured using a BCA protein assay kit (Pierce Biotechnology Inc., Rockford, IL, USA).

For immunoblotting, equal amounts of protein (65 μ g) were loaded into each well of a 7.5% Tris-glycine pre-cast polyacrylamide gel (Bio-Rad Laboratories, Hercules, CA, USA) and separated by electrophoresis. After separation, the proteins were transferred from the gel to nitrocellulose membrane (Bio-Rad Laboratories). The nitrocellulose membrane was then incubated overnight at 4°C in primary antibody solution, namely c-Jun (1:500 dilution, polyclonal c-Jun antibody; Cell Signaling Technology Inc., Danvers, MA, USA), c-Fos (1:500 dilution, polyclonal c-Fos antibody; Cell Signaling Technology Inc.), or histone H3 (as an internal standard, 1:500 dilution, polyclonal histone H3 antibody; Cell Signaling Technology Inc.).

Horseradish peroxidase conjugated anti-mouse IgG antibody (Amersham Pharmacia Biotech Inc.,

Piscataway, NJ, USA) was next used as the secondary antibody. Bound antibody was detected by chemiluminescence (ECL plus kit; Amersham Pharmacia Biotech Inc.) and high performance film (Hyperfilm; Amersham Pharmacia Biotech Inc.). The protein band densities were quantified using densitometry (Scion Corp., Frederick, MD, USA).

2.4. Immunofluorescent staining for AP-1

We chose to employ immunofluorescent staining to elucidate the effect of MgSO_4 on AP-1 nuclear translocation. To facilitate immunofluorescent staining, RAW264.7 cells were treated with PBS, MgSO_4 , LPS, LPS plus MgSO_4 , or LPS plus MgSO_4 plus BAY-K8644 and harvested at 1 hour after the start of treatment. The timings for cell harvesting were based on the preliminary c-Jun data obtained from the immunoblotting assays. After harvesting, immunofluorescent staining was performed as previously described [14]. In brief, RAW264.7 cells were fixed with freshly prepared 3% formaldehyde at room temperature for 5 minutes and permeabilized by 0.5% Triton X-100 (Sigma-Aldrich) on ice for 5 minutes. Cells were then rinsed twice with $1\times$ PBS. After 30 minutes of blocking with 1% bovine serum albumin (Sigma-Aldrich), the RAW264.7 cells were then incubated in c-Jun primary antibody solution (1:100 dilution, polyclonal c-Jun antibody; Cell Signaling Technology Inc.) for 30 minutes followed by incubation with fluorescent rhodamine isothiocyanate-conjugated secondary antibody (Jackson ImmunoResearch Inc., West Grove, PA, USA) for 30 minutes. Nuclei were counterstained with diamidino-2-phenylindole (Pierce Biotechnology Inc.) and the cells were viewed using a microscope.

2.5. Electrophoretic mobility shift assay (EMSA)

After the RAW264.7 cells had been treated with PBS, MgSO_4 , LPS, LPS plus MgSO_4 , or LPS plus MgSO_4 plus BAY-K8644, they were harvested 1 hour after treatment began in order to facilitate EMSA analysis of AP-1. After harvesting, nuclear extracts were prepared, as described above. EMSA was then performed using a chemiluminescence EMSA kit (AP-1 EMSA kit, respectively; Panomics Inc., Fremont, CA, USA), as previously described [15].

2.6. Statistical analysis

One way analysis of variance was performed to test the between-group differences. The Tukey test was used for multiple comparisons. Data are presented

as mean \pm standard deviation. The significance level was set at 0.05. A commercial software package (SPSS 11.5 for Windows; SPSS Inc., Chicago, IL, USA) was used for data analyses.

3. Results

3.1. MgSO_4 inhibited LPS-induced AP-1 activation

PBS or MgSO_4 had no significant effect on AP-1 expression, based on the fact that the c-Jun and c-Fos protein concentrations from the PBS and M groups were low throughout the experiments (data not shown). In contrast, the c-Jun and c-Fos protein concentrations of the LPS group, from cell cultures harvested at 1, 3, and 5 hours after LPS administration, were significantly higher than those of the PBS group (all $p < 0.001$, data not shown). Moreover, the c-Jun protein concentrations of the LPS+M group, from cell cultures harvested at 1, 3, and 5 hours after LPS administration, were significantly lower than those of the LPS group ($p = 0.023$, 0.025, and 0.011, respectively; Fig. 1). The c-Fos protein concentrations of the LPS+M group, from cell cultures harvested at 1 and 3 hours after LPS administration, were also significantly lower than those of the LPS group ($p = 0.005$ and 0.001, respectively; Fig. 1).

AP-1 nuclear translocation was evaluated by immunofluorescent staining. The immunofluorescent staining results confirmed that LPS induced significant AP-1 nuclear translocation, as the fluorescence intensity of the LPS group was stronger than that of the PBS group (data not shown). Moreover, MgSO_4 significantly inhibited the effect of LPS on the induction of AP-1 nuclear translocation as the fluorescence intensity of the LPS+M group was weaker than that of the LPS group (Fig. 2).

The AP-1 DNA-binding activity of the PBS group was low (Fig. 3A). The AP-1 DNA-binding activity of the M group was comparable to that of the PBS group, whereas the AP-1 DNA-binding activity of the LPS group was significantly higher than that of the PBS group ($p < 0.001$; Fig. 3A). Moreover, the AP-1 DNA-binding activity of the LPS+M group was significantly lower than that of the LPS group ($p = 0.021$; Fig. 3).

3.2. Effects of MgSO_4 on AP-1 were reversed by BAY-K8644

The c-Jun protein concentrations of the LPS+M+K group in cell cultures harvested at 1 and 5 hours after LPS administration were significantly higher than those of the LPS+M group ($p = 0.036$ and 0.019, respectively; Fig. 1). The c-Fos protein concentrations




Fig. 1 — Representative gel photography and densitometric analysis of the nuclear protein concentrations of activator protein-1 (AP-1), including c-Jun and c-Fos, in the nuclear extracts of RAW264.7 cells after treatment with lipopolysaccharide (LPS), LPS plus magnesium sulfate (MgSO_4), and LPS plus MgSO_4 plus the potent L-type calcium channel activator BAY-K8644 (denoted as the LPS, LPS+M, and LPS+M+K groups, respectively). The c-Jun and c-Fos concentrations were normalized against histone H3. Data are expressed as mean \pm standard deviation. * $p < 0.05$, LPS+M group vs. LPS group; † $p < 0.05$, LPS+M+K group vs. LPS+M group.

of the LPS+M+K group in cell cultures harvested at 1, 3, and 5 hours after LPS administration were also significantly higher than those of the LPS+M group ($p = 0.017$, 0.033 , and 0.025 , respectively; Fig. 1). The fluorescence intensity of the LPS+M+K group was stronger than that of the LPS+M group (Fig. 2). Moreover, the AP-1 DNA-binding activity of the LPS+M+K group was significantly higher than that of the LPS+M group ($p = 0.026$; Fig. 3).

4. Discussion

The results from this study confirm our hypothesis that MgSO_4 is able to attenuate AP-1 activation in endotoxin-activated murine macrophages. The crucial role of AP-1 in the upregulation of inflammatory molecules during sepsis is well-established (4). Our results, in concert with previous ones (9,10), confirm the potent anti-inflammation capacity of MgSO_4 . In addition, these results provide clear evidence to support the notion that MgSO_4 , in order to exhibit its anti-inflammation effects, may act by inhibiting endotoxin-induced AP-1 activation.

Endotoxin induces a transient increase in intracellular calcium level due to an initial intracellular calcium

release followed by calcium influx from the extracellular space (16). This endotoxin-induced transient increase in intracellular calcium plays a crucial role in regulating the expression of the signaling pathways and subsequent inflammatory molecules, as expression of these signaling pathways and the inflammatory molecules are inhibited by calcium chelators (16). The L-type calcium channels are members of a transmembrane ion channel group of proteins that are associated with calcium influx signaling in immune cells (17). As mentioned earlier, magnesium is a natural calcium antagonist and a potent calcium channel inhibitor (11). The results from this study reveal that the effects of MgSO_4 on inhibiting AP-1 activation are significantly reversed by the L-type calcium channel activator BAY-K8644. This supports the notion that MgSO_4 may act by inhibiting calcium influx from the extracellular space via the L-type calcium channels and that this leads to its inhibition of AP-1 activation.

Cellular recognition of endotoxin, including the binding of endotoxin to inflammatory cells and subsequent toll-like receptor 4 (TLR-4) activation, has been shown to play an essential role in mediating endotoxin-induced activation of AP-1 (18). Cellular recognition of endotoxin also involves the membrane-anchor protein complex cluster of differentiation-14




Fig. 2 — Representative immunofluorescent staining findings for activator protein-1 (AP-1) nuclear translocation of RAW264.7 cells after treatment with lipopolysaccharide (LPS), LPS plus magnesium sulfate (MgSO_4), and LPS plus MgSO_4 plus the potent L-type calcium channel activator BAY-K8644 (denoted as the LPS, LPS+M, and LPS+M+K groups, respectively). AP-1 (c-Jun) was stained with fluorescent rhodamine isothiocyanate-conjugated antibody. The nuclei were counterstained with diaminido-2-phenylindole. The cells were imaged using a fluorescence microscope.




Fig. 3 — (A) Representative gels showing activator protein-1 (AP-1) DNA-binding activity in the nuclear extracts of RAW264.7 cells after treatment with phosphate buffered saline (PBS), lipopolysaccharide (LPS), LPS plus magnesium sulfate (MgSO_4), and LPS plus MgSO_4 plus the potent L-type calcium channel activator BAY-K8644 (denoted as the PBS, LPS, LPS+M, and LPS+M+K groups, respectively) by chemiluminescence electrophoretic mobility shift assay. (B) Densitometric analyses of the band intensities of the LPS, LPS+M and LPS+M+K groups. * $p < 0.05$, LPS+M group vs. LPS group; † $p < 0.05$, LPS+M+K group vs. LPS+M group.

(CD14) [18,19]. After binding, endotoxin induces a physical proximity of CD14 and TLR-4 that allows CD14 to present endotoxin to TLR-4, which then leads to the activation of intracellular signaling pathways [18,19]. Direct evidence delineating the effect of MgSO_4 on the regulation of endotoxin binding to inflammatory cells and endotoxin-induced TLR-4 and CD14 activation is still lacking. Magnesium is an antagonist of the N-methyl-D-aspartate (NMDA) receptors [20]. Previous findings have indicated that inhibition of the NMDA receptors may result in an attenuation of TLR-4 activation [21]. Other results have indicated that rats fed a magnesium-deficient diet show enhanced CD14 expression in their intestinal and cardiac tissues [22]. These results seem to suggest the possibility that endotoxin-induced TLR-4 and/or CD14 activation may be regulated by MgSO_4 . In addition, myeloid differential-2 (MD-2), an extracellular protein associated with the extracellular domain of TLR-4, is also required for the activation of TLR-4 by endotoxin [18]. MD-2 comprises a cationic and a hydrophobic binding site [23]. As magnesium is a divalent cation, we speculate that endotoxin-induced MD-2 activation might also be regulated by MgSO_4 . A follow-up study is currently being conducted in our laboratory to elucidate the effects of MgSO_4 on the regulation of endotoxin-inflammatory cells binding and the endotoxin-induced TLR-4/CD14/MD-2 complex.

Certain limitations to this study exist. First, although the findings of this study are in agreement with those of previous studies [9,10] and confirm the potent anti-inflammatory capacity of MgSO_4 , the effects of MgSO_4 in terms of protecting vital organs from the detrimental effects of endotoxin remain unstudied. Second, while our results confirm the involvement of L-type calcium channels in mediating the anti-inflammatory effects of MgSO_4 , this leaves unanswered the potential roles of the other calcium channels, which remain to be determined. Third, the question of whether the anti-inflammatory effects of MgSO_4 are also involved in other crucial mechanisms, such as the NMDA receptors, remains unanswered. Fourth, it is important to bear in mind that interspecies differences need to be considered when extending the interpretation of the present results.

In conclusion, MgSO_4 inhibited AP-1 activation in LPS-treated macrophages and the mechanisms would seem to involve the L-type calcium channels.

Acknowledgments

This work was supported by grants from Buddhist Tzu Chi General Hospital, Taipei Branch (TCRD-TPE-99-34) awarded to C.H. Chen and the National Science Council, Taiwan (NSC 98-2314-B-303-012-MY3) awarded to C.J. Huang.

References

- Blackwell TS, Christman JW. Sepsis and cytokines: current status. *Br J Anaesth* 1996;77:110–7.
- Yoshinari D, Takeyoshi I, Koibuchi Y, et al. Effects of a dual inhibitor of tumor necrosis factor- α and interleukin-1 on lipopolysaccharide-induced lung injury in rats: involvement of the p38 mitogen-activated protein kinase pathway. *Crit Care Med* 2001;29:628–34.
- Blackwell TS, Christman JW. The role of nuclear factor- κ B in cytokine gene regulation. *Am J Respir Cell Mol Biol* 1997;17:3–9.
- Adcock IM. Transcription factors as activators of gene transcription: AP-1 and NF- κ B. *Monaldi Arch Chest Dis* 1997;52:178–86.
- Kerr LD, Inoue J, Verma IM. Signal transduction: the nuclear target. *Curr Opin Cell Biol* 1992;4:496–501.
- Bar J, Ben Haroush A, Feldberg D, Hod M. The pharmacologic approach to the prevention of preeclampsia: from antiplatelet, antithrombosis and antioxidant therapy to anticonvulsants. *Curr Med Chem Cardiovasc Hematol Agents* 2005;3:181–5.
- Telci L, Esen F, Akcora D, Erden T, Canbolat AT, Akpir K. Evaluation of effects of magnesium sulphate in reducing intraoperative anaesthetic requirements. *Br J Anaesth* 2006; 89:594–8.
- Brill S, Sedgwick PM, Hamann W, Di Vadi PP. Efficacy of intravenous magnesium in neuropathic pain. *Br J Anaesth* 2002;89:711–4.
- Lin CY, Tsai PS, Hung YC, Huang CJ. L-type calcium channels are involved in mediating the anti-inflammatory effects of magnesium sulphate. *Br J Anaesth* 2010;104: 44–51.
- Rochelson B, Dowling O, Schwartz N, Metz CN. Magnesium sulfate suppresses inflammatory responses by human umbilical vein endothelial cells (HuVECs) through the NF κ B pathway. *J Reprod Immunol* 2007;73:101–7.
- Sonna LA, Hirshman CA, Croxton TL. Role of calcium channel blockade in relaxation of tracheal smooth muscle by extracellular Mg^{2+} . *Am J Physiol* 1996;271:L251–7.
- Jones E, Adcock IM, Ahmed BY, Punchard NA. Modulation of LPS stimulated NF- κ B mediated nitric oxide production by PKC ϵ and JAK2 in RAW macrophages. *J Inflamm (Lond)* 2007;4:23.
- Tsai PS, Chen CC, Tsai PS, Yang LC, Huang WY, Huang CJ. Heme oxygenase 1, nuclear factor E2-related factor 2, and nuclear factor κ B are involved in hemin inhibition of type 2 cationic amino acid transporter expression and L-arginine transport in stimulated macrophages. *Anesthesiology* 2006;105:1201–10.
- Park GY, Le S, Park KH, et al. Anti-inflammatory effect of adenovirus-mediated IkappaB α overexpression in respiratory epithelial cells. *Eur Respir J* 2001;18:801–9.
- Nareika A, He L, Game BA, et al. Sodium lactate increases LPS-stimulated MMP and cytokine expression in U937 histiocytes by enhancing AP-1 and NF- κ B transcriptional activities. *Am J Physiol Endocrinol Metab* 2005;289: E534–42.
- Zhou X, Yang W, Li J. Ca^{2+} - and protein kinase C-dependent signaling pathway for nuclear factor- κ B activation, inducible nitric-oxide synthase expression, and tumor necrosis factor- α production in lipopolysaccharide-stimulated rat peritoneal macrophages. *J Biol Chem* 2006;281: 31337–47.

17. Azenabor AA, Chaudhry AU. Effective macrophage redox defense against *Chlamydia pneumoniae* depends on L-type Ca^{2+} channel activation. *Med Microbiol Immunol* 2003;192:99–106.
18. Heumann D, Roger T. Initial responses to endotoxins and Gram-negative bacteria. *Clin Chim Acta* 2002;323:59–72.
19. Jiang Q, Akashi S, Miyake K, Petty HR. Lipopolysaccharide induces physical proximity between CD14 and toll-like receptor 4 (TLR4) prior to nuclear translocation of NF-kappa B. *J Immunol* 2000;165:3541–4.
20. Lodge D, Johnson KM. Noncompetitive excitatory amino acid receptor antagonists. *Trends Pharmacol Sci* 1990;11:81–6.
21. Wu GJ, Chen TL, Ueng YF, Chen RM. Ketamine inhibits tumor necrosis factor-alpha and interleukin-6 gene expressions in lipopolysaccharide-stimulated macrophages through suppression of toll-like receptor 4-mediated c-Jun N-terminal kinase phosphorylation and activator protein-1 activation. *Toxicol Appl Pharmacol* 2008;228:105–13.
22. Chmielinska JJ, Tejero-Taldo MI, Mak IT, Weglicki WB. Intestinal and cardiac inflammatory response shows enhanced endotoxin receptor (CD14) expression in magnesium deficiency. *Mol Cell Biochem* 2005;278:53–7.
23. Jerala R. Structural biology of the LPS recognition. *Int J Med Microbiol* 2007;297:353–63.